



# LIQUIDS PRODUCTS

P

- 2 AD3E+CK
- 3 ALAN AD3EC
- 4 ALAN AD3E FORT
- 5 ALAN AD3E
- 6 SELEVIT-E20
- 7 SELEVIT-E
- 8 AMINO+B
- 9 VITALAN

P

- 10 SUPER-VITALAN
- 11 MINALAN SOL
- 12 ALAN OIL
- 13 HI-IMMUNE
- 14 SUPER-IMMUNE
- 15 HEPALAN
- 16 RENALAN
- 17 BRONCHI OIL

P

- 18 ELECTROALAN
- 19 KETOLAN
- 20 ALANTOX
- 21 TOXANIL

# AD3E+CK



## COMPOSITION:

Each 1000 ml contains:

Vitamin A	40,000,000	IU
Vitamin D3	5,000,000	IU
Vitamin E	50,000	mg
Vitamin C	25,000	mg
Vitamin K3	25,000	mg

## INDICATIONS:

- Increase animal vitality and maintain good health.
- Support productivity.
- Support and formation of vital immunity.
- Relieving symptoms of heat stress.

## DOSAGE:

- **Poultry:** 50 ml per 200 litres of drinking water from 3-5 day.
- **Cattle, Horses:** 1 ml per 10 Kg bodyweight for 3-5 day.
- **Sheep, Goats, Calves:** 1 ml per 10 Kg bodyweight for 3-5 day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# ALAN AD3EC



## COMPOSITION:

Each 1000 ml contains:


Vitamin A	50,000,000	IU
Vitamin D3	2,500,000	IU
Vitamin E	25,000	mg
Vitamin C	6,000	mg

## INDICATIONS:

- Increase animal vitality and maintain good health.
- Support productivity.
- Support and formation of vital immunity.
- Relieving symptoms of heat stress.

## DOSAGE:

- **Poultry:** 50 ml per 200 litres of drinking water from 3-5 day.
- **Cattle:** 1000 ml per 1000 litres of drinking water from 3-5 day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.



# ALAN AD3E FORT



## COMPOSITION:

Each 1000 ml contains:


Vitamin A	25,000,000	IU
Vitamin D3	7,000,000	IU
Vitamin E	25,000	mg

## INDICATIONS:

- Increase animal vitality and maintain good health.
- Support productivity.
- Support and formation of vital immunity.
- Relieving symptoms of heat stress.

## DOSAGE:

- **Poultry:** 50ml per 200 litres of drinking water from 3-5 day.
- **Cattle, Horses:** 5 - 10 ml per animal during 3 - 5 day.
- **Sheep, Goats, Calves:** 2 - 3 ml per animal during 3 - 5 day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# ALAN AD3E



## COMPOSITION:

Each 1000 ml contains:

Vitamin A	10,000,000	IU
Vitamin D3	5,000,000	IU
Vitamin E	5,000	mg

## INDICATIONS:

Increase animal vitality and maintain good health.

Support productivity.

Support and formation of vital immunity.

## DOSAGE:

- **Poultry:** 100 per 200 litres of drinking water from 3-5 day.
- **Cattle, Horses:** 10 - 15 ml per animal during 2 - 3 day.
- **Foals, Calves:** 15 ml per animal during 2 - 3 day.
- **Sheep, Goats:** 10 - 15 ml per animal during 2 - 3 day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# SELEVIT-E20



## COMPOSITION:

Each 1000 ml contains:

Vitamin E	200,000	mg
Selenium	900	mg

## INDICATIONS:

- Compensation for the deficiency of this vitamin in the animal's body.
- Protection of chicks from encephalopathy (crazy chick disease).
- Prevention of muscular dystrophy (white muscle disease, rigors of pregnancy).
- Increased fertility and production.

## DOSAGE:

- Calves: 1 – 3 ml / 1 litter for 3-5 day.
- Cows: 7.5 - 15 ml / 1 litter for 3-5 day.
- Sheep: 2 - 5 ml / 1 litter for 3-5 day.
- Lambs: 1 – 3 ml / 1 litter for 3-5 day.
- Poultry: 50 ml / 200 litter of drinking water from 3-5 day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.



# SELEVIT-E



## COMPOSITION:

Each 1000 ml contains:

Vitamin E	100,000	mg
Selenium	450	mg

## INDICATIONS:

- Compensation for the deficiency of this vitamin in the animal's body.
- Protection of chicks from encephalopathy (crazy chick disease).
- Prevention of muscular dystrophy (white muscle disease, rigors of pregnancy).
- Increased fertility and production.

## DOSAGE:

- Calves: 3 - 5 ml / 1 litter for 3-5 day.
- Cows: 20 - 25 ml / 1 litter for 3-5 day.
- Sheep: 5 - 10 ml / 1 litter for 3-5 day.
- Lambs: 2 - 5 ml / 1 litter for 3-5 day.
- Poultry: 100 ml / 200 litter of drinking water from 3-5 day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# AMINO+B



## COMPOSITION:

Each 1000 ml contains:


Vitamin B2	215 mg
Vitamin B3 (Niacin)	3,500 mg
Vitamin B5	135 mg
Vitamin B6	125 mg
Vitamin B12	1,250 mcg
DL-Methionine	1,000 mg
Choline Chloride	750 mg
L-Lysine Hcl	1,000 mg

## INDICATIONS:

- Increasing the rate of feed consumption.
- Supporting the activity of the digestive system.
- Increase conversion rate.
- Supplying the animal body with the necessary amino acids.

## DOSAGE:

- Poultry: 100 ml / 100 L of drinking water.
- Calves and Lambs: 10 ml / 50 kg body weight.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.



# VITALAN



## COMPOSITION:

Each 1000 ml contains:

Vitamin A	12,000,000	IU	L-Threonine	441	mg
Vitamin D3	500,000	IU	L-Valin	828	mg
Vitamin E	10,000	mg	L-Alanine	585	mg
Vitamin B1	3,000	mg	L-Arginine	729	mg
Vitamin B2	500	mg	Aspartic Acid	630	mg
Niacin	10,000	mg	L-Phenylalanine	540	mg
Calcium- D-Pantothenate	4,000	mg	Glycine	927	mg
Vitamin B6	1,000	mg	Monosodium Glutamate	1,800	mg
Biotin	15,000	mcg	L-Isoleucine	540	mg
Vitamin B12	20,000	mcg	L-Leucine	486	mg
Vitamin C	7,000	mg	L-Lysine Hcl	1,557	mg
Choline Chloride	8,000	mg	L-Proline	1,116	mg
Carnitine	1,500	mg	L-Tyrosine	291.6	mg
DI-Methionine	340.2	mg			

## INDICATIONS:

Increase animal vitality and maintain its health.  
Productivity support.  
Support and formation of vital immunity.  
Relieving symptoms of heat stress.

Providing the animal body with an integrated set of amino acids that play an important role in most metabolic processes within the body.

## DOSAGE:

- **Poultry:** 100 ml / 200 L of drinking water for 3 – 5 days.
- **Calves and lambs:** 1 ml / 10 kg body weight daily.
- **Horses:** 10 ml / 100 kg body weight 3-5 days before the race.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# SUPER-VITALAN



## COMPOSITION:

Each 1000 ml contains:

Vitamin A	8,000,000 IU	Iodate	10 mg
Vitamin D3	1,000,000 IU	Potassium	900 mg
Vitamin E	10,000 mg	Selenium	200 mg
Vitamin B1	4,000 mg	Methionine	945 mg
Vitamin B2	1,000 mg	Threonine	1,225 mg
Niacin	6,000 mg	Valin	2,300 mg
Choline Chloride	6,500 mg	Alanine	1,625 mg
Pantothenic acid	4,000 mg	Arginine	2,025 mg
Vitamin B6	1,000 mg	Aspartic acid	1,750 mg
Vitamin B7	15 mg	Phenylalanine	1,500 mg
Vitamin B12	5 mg	Glycine	2,575 mg
Vitamin C	900 mg	Glutamic acid	3,525 mg
Carnitine	1,200 mg	Isoleucine	1,360 mg
Iron	1,800 mg	Lucien	2,425 mg
Zinc	4,000 mg	Lysine	4,325 mg
Copper	100 mg	Proline	3,100 mg
Manganese	230 mg	Tyrosine	810 mg
Phosphorus	2,400 mg	Aroma	10,000 mg
Cobalt	50 mg		

## INDICATIONS:

It is used as a Supplementary Ground in order to meet the vitamin, amino acid, trace mineral and aromatic substance needs of Adult Rams, Sheep, Goats, Tosun, Dairy Cows and Beef Cattles, and Lambs, Capricorns and Calves that have completed their rumen development.

## DOSAGE:

- Adult Rams and Goats 10-12 ml per animal per day.
- Adult Sheep and Goats 6-7 ml per animal per day.
- 3-4 ml per animal per day for Lambs and Kids that have completed their rumen development.
- Adult Tosun Dairy Cows and Fattening Cattles 10-12 ml per 100 kg body weight per day.
- It is used by mixing 6-7 ml of drinking water per animal per day for calves that have completed their rumen development.
- Poultry: 100 ml / 200 L of drinking water for 3 – 5 days

## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine



IT'S USED FOR:



MEDICATED DRINKING WATER SHOULD BE USED WITHIN 24 HOURS.



# MINALAN SOL



## COMPOSITION:

Each 1000 ml contains:

Iron	25,000	mg
Zinc	8,000	mg
Phosphor	4,900	mg
Cobalt	350	mg
Copper	5,850	mg

## INDICATIONS:

Supplying the herd with a group of minerals necessary to compensate for the deficiency.

Activate the intestines and increase the rate of absorption.

Herd immunity support.

## DOSAGE:

- **Poultry:** 200 ml / 200 L of drinking water.
- **Calves and lambs:** 5 ml / 50 kg body weight.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.



# ALAN OIL



## COMPOSITION:

Each 1000 ml contains:

Garlic Oil	50,000 mg
Thyme Oil	10,000 mg
Oregano Oil	25,000 mg

## INDICATIONS:

Immune system support.

Parasite control.

Reducing the severity of symptoms in bacterial infections.

Protect and increase liver activity and bile secretion.

## DOSAGE:

- Poultry: 100 ml per 200 L of drinking water.
- Cattle, Horses: 1 ml per 10 kg bodyweight for 3 – 5 days.
- Sheep, Goats, Calves: 1 ml per 10 kg bodyweight for 3 – 5 days.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# HI-IMMUNE



## COMPOSITION:

Each 1000 ml contains:

Yeast Extract	80,000 mg
Artichoke Extract	20,000 mg
Citrus Extract	20,000 mg
Sorbitol	200,000 mg

## INDICATIONS:

Supporting the liver and increasing its functional activity.  
Supporting the digestive system and stimulating digestion.  
Increase production rate. Supporting the immune system in birds.

## DOSAGE:

- **Poultry:** 500 ml per 1000 L of drinking water.
- **Cattle, Horses:** 1 ml per 10 kg bodyweight for 3 – 5 days.
- **Sheep, Goats, Calves:** 1 ml per 10 kg bodyweight for 3 – 5 days.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# SUPER-IMMUNE



## COMPOSITION:

Each 1000 ml contains:

L-Lysine Hcl	30,000 mg
L-Threonine	45,000 mg
DL-Methionine	6,000 mg
Citric Acid	68,600 mg
Lactic Acid	88,000 mg
Zinc Glycinate	60,000 mg
Copper Glycinate	30,000 mg
Yeast Extracte	80,000 mg

## INDICATIONS:

- Supporting the immune system in birds.
- Supporting the liver and increasing its functional activity.
- Supports the digestive system and stimulates digestion.
- Increase production rate.

## DOSAGE:

- Poultry: 1 ml per 3 litres of water.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED FOR:



MEDICATED DRINKING WATER SHOULD BE USED WITHIN 24 HOURS.



# HEPALAN



## COMPOSITION:

Each 1000 ml contains:

Silymarin	100,000	mg
Sorbitol	100,000	mg
L-Carnitine	25,000	mg
Choline Chloride	25,000	mg
Propionic Acid	100,000	mg
Nicotinamide	50,000	mg
Lactic Acid	50,000	mg
DL-Methionine	4,000	mg
Betaine	5,000	gm

## DOSAGE:

For oral administration via feed or drinking water.

- **Poultry:** 50-100 ml per 200 liter of drinking water for 3-5 days.
- **Cattle, Cow:** 5 ml / 100 kg B.W for 3-5 days
- **Sheep, Goats, Calves:** 5 ml / 50 kg B.W for 3-5 days
- **Horse, Foal:** 7.5 ml / 100 kg B.W for 3 – 5 days

## INDICATIONS:

For the proper function of liver functions: metabolism, detoxification, secretion, and productivity. Protects the liver against toxic effects.

For B-Complex vitamin deficiencies and retarded growth.

Increase weight gain and stimulate the metabolism. Prevents fatty liver syndrome and improves liver functions.

Improves carbohydrate metabolism for proper functioning of nervous tissue, cardiac muscle and enzyme activities.

Important role in metabolism, growth rate and feed utilization.

Stimulates regeneration of hepatic cells and better productivity.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED FOR:



MEDICATED DRINKING WATER SHOULD BE USED WITHIN 24 HOURS.

# RENALAN



## COMPOSITION:

Each 1000 ml contains:

Sorbitol	50,000	mg
Ammonium Chloride	300,000	mg
Magnesium Sulphate	10,000	mg
DL-Methionine	48,000	mg

## INDICATIONS:

- Activates kidney functions
- Prevents fluid retention inside the body
- Protecting the kidneys and liver from toxins.

## DOSAGE:

For oral administration via feed or drinking water.

- **Poultry:** 75 ml per 200 litters of drinking water for 3 – 5 days.
- **Cattle, Horses:** 10 ml per 100 kg bodyweight for 3 – 5 days.
- **Sheep, Goats, Calves:** 20 ml per 100 kg bodyweight for 3 – 5 days.
- Mixed with feed, the product should be used immediately.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# BRONCHI OIL



## COMPOSITION:

Each 1000 ml contains:

Eucalyptus Oil	20,000	mg
Thyme Oil	10,000	mg
Menthol	25,000	mg

## INDICATIONS:

Expectorant and bronchodilator.

Reducing the severity of respiratory symptoms.

Appetizing.

## DOSAGE:

- **Poultry:** 500 ml per 1000 litters of drinking water.
- **Cattle, Horses:** 1 ml per 10 kg bodyweight for 3 – 5 days.
- **Sheep, Goats, Calves:** 1 ml per 10 kg bodyweight for 3 – 5 days.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.



# ELECTROALAN



## COMPOSITION:

Each 1000 ml contains:

Calcium Chloride	97,107	mg
Magnesium Chloride	38,681	mg
Sodium Chloride	5,080	mg
Potassium Chloride	1,906	mg
Iron Chloride	4,357	mg
Cobalt Chloride Hexahydrate	404	mg
Zinc Chloride	8,336	mg
Copper Chloride Pentahydrate	2,684	mg
Sodium Selenate 45%	239	mg
Phosphoric Acid	134,470	mg
DL-Methionine	1,000	mg
L-Lysine Hcl	1,000	mg

## INDICATIONS:

- Treatment of metabolic disorders.
- Wasting treatment.
- Mineral deficiency.
- Increase production and eggshell hardness in poultry.

## DOSAGE:

- **Poultry:** 300 ml / 1000 L of drinking water for 3-5 days.
- **Calves and Lambs:** 1 ml / 20 kg body weight 3-5 days.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# KETOLAN



## COMPOSITION:

Each 1000 ml contains:

Propylene Glycol	300,000	mg
Glycerine	665,000	mg
Vitamin A	300,000	IU
Vitamin D3	75,000	IU
Vitamin E	300	mg
Niacin	300	mg
Carnitine	5,000	mg
L-Lysine Hcl	43.25	mg
DI-Methionine	9.45	mg
L-Threonine	12.25	mg
Monosodium Glutamate	50	mg

L-Proline	31	mg
L-Glycine	25.75	mg
L-Leucine	24.25	mg
L-Valin	23	mg
L-Arginine	20.25	mg
Aspartic Acid	17.5	mg
L-Alanine	16.25	mg
L-Phenylalanine	15	mg
L-Isoleucine	13.5	mg
L-Tyrosine	8.1	mg

## INDICATIONS:

- Prevention of possible diseases during childbirth.
- A growth and performance booster.
- Increase feed conversion rate.
- Immune system support.

## DOSAGE:

- 300 - 500 ml/day from 3 days before birth to 4 days after birth.
- Used orally.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.

# ALANTOX



## COMPOSITION:

Each 1000 ml contains:

Phosphoric Acid	57,800 mg	Acetic Acid	17,600 mg
Citric Acid	90,000 mg	Betaine HCL	12,000 mg
Lactic Acid	48,000 mg	Copper Sulphate	12,000 mg
Malic Acid	25,000 mg	L-Lysine Hcl	18,000 mg
Tartaric Acid	12,000 mg		
Vitamin B6	2,100 mg		
Propylene Glycol	60,000 mg		
Yeast Extract	45,000 mg		
Formic Acid	8,500 mg		
Propionic Acid	30,000 mg		

## INDICATIONS:

- A growth and performance booster.
- Increase feed conversion rate.
- Immune system support.
- Prevention of bacterial infections.
- Protection from infection with mycotoxins

## DOSAGE:

- **Poultry:** 100 ml per 200 litters of water.
- **Cows and Sheep:** 1 ml / 10 kg live weight per day.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.



# TOXANIL



## COMPOSITION:

Each 1000 ml contains:

Citric Acid	25,000 mg
Lactic Acid	11,000 mg
Sorbic Acid	3,000 mg
Malic Acid	21,000 mg
Acetic Acid	70,000 mg
Propionic Acid	160,000 mg
Phosphoric Acid	22,000 mg
Formic Acid	75,000 mg

## INDICATIONS:

- Control and reduce feed contaminations.
- Bind multiple mycotoxins, inhibits growth of mold and prevents further mycotoxin production.
- Increases immunity and reduce stress on liver.
- Improves laying performance and growth in birds.
- Reduces pH in gastro-intestinal tract.
- Reliable protection from spoilage.
- Targetted reduction of harmful bacteria and fungus.
- Long term control of Salmonella, E.coli and other harmful bacteria.
- Support of optimum performance

## DOSAGE:

- 100 ml per 200 litters of drinking water for 3-5 days.



## SUPPLEMENT

It is a Feed Additive  
Not Used as Medicine

IT'S USED  
FOR:



MEDICATED DRINKING WATER SHOULD  
BE USED WITHIN 24 HOURS.